PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6:

C07K 5/02, A61K 38/05

(11) International Publication Number:

WO 98/09986

A1

(43) International Publication Date:

12 March 1998 (12.03.98)

(21) International Application Number:

PCT/GB97/02358

(22) International Filing Date:

1 September 1997 (01.09.97)

(30) Priority Data:

9618277.9

2 September 1996 (02.09.96) GB

9700896.5 17 January 1997 (17.01.97) GB

(71) Applicant (for all designated States except US): THE MAN-CHESTER METROPOLITAN UNIVERSITY [GB/GB]; All Saints Building, All Saints, Manchester M15 6BH

(72) Inventor; and

- (75) Inventor/Applicant (for US only): D'SILVA, Claudius [GB/GB]; 23 Chiltern Drive, Hale/Altringham, Cheshire WA15 9PL (GB).
- (74) Agents: EVANS, David, Charles et al.; F.J. Cleveland & Company, 40-43 Chancery Lane, London WC2A 1JQ (GB).

(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).

Published

With international search report.

(54) Title: S-BLOCKED GLUTATHIONES

(57) Abstract

This invention concerns glutathione derivatives, and in particular their application in the suppression of pathogens. It has been discovered that certain glutathione derivatives are effective inhibitors of the growth of a range of cancer cell types, and certain microorganisms. According to one aspect of the present invention there is provided a glutathione having structure (I). Compounds based upon this general structure are disclosed which are active against parasitic infectious agents such as T. Brucei and L. Donovani. Further compounds are disclosed which are active against cancer cells.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	Si	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT.	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CF	_	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Congo Switzerland	KG	Kyrgyzstan	NO	Norway	zw	Zimbabwe
	•	KP	Democratic People's	NZ	New Zealand		
Cl	Côte d'Ivoire	***	Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Ponugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
cz	Czech Republic	น	Liechtenstein	SD	Sudan		
DE	Germany		Sri Lanka	SE	Sweden		
DK	Denmark	LK LR	Liberia	SG	Singapore		
EE	Estonia	LK	Liucha	30	Sin-Supo. o		

S-Blocked Glutathiones

This invention concerns glutathione derivatives, and
in particular their application in the suppression of pathogens.

Illness may be caused by many agents. Bacterial infections are caused by micro-organisms which

multiply rapidly and cause a number of diseases.

Parasites are organisms which live in or on a host and feed off the host. Cancers are evident by the uncontrolled multiplication of cells in the body. The cancer may be localized, such as breast cancer, or systemic such as leukaemia.

The treatment of the illnesses caused by the aforementioned agents has been the subject of much research, and many different approaches. One approach involves targeting the cells which cause the disease and destroying them or disrupting their ability to multiply.

For such an approach to be successful the agent or

drug used to attack the diseased cells should not harm
significantly other healthy cells. Thus such an

approach requires an understanding and identification of the biochemical processes carried out in cells, and the targeting and disruption of specific processes which are unique to the diseased cells.

5

Such an approach involves the use of the cytotoxic compound methylgloxal which is produced in the cells of certain organisms. A build up of methylgloxal in the cell up to cytotoxic levels will, of course, result in cell death. By inducing a build up of methylgloxal, significant growth inhibition effects have been seen in tumour cells, Escherichia coli, Saccharomycescerevisise and Leishmania donovani.

The build up of methylgloxal may be promoted by the inhibition of glyoxalase I (GLI) enzyme. A wide range of inhibitors of GLI are known and these include substrate or product analogues and mechanism-based inhibitors.

20

25

Glutathione (\gamma-glutamylcysteinylglycine) fulfils a variety of roles vital to life processes. It functions as a co-enzyme, co-substrate, substrate or part of the substrate architecture. S-blocked glutathiones have been shown to be potent inhibitors of GLI in vitro and this has led to a search for particular S-blocked

15

glutathiones which may have therapeutic effect.

A general procedure for the preparation of the monoglycyl and dimethyl ester and amide derivatives of \underline{S} -(4-bromobenzyl)glutathione has been described by the inventor in Biochem. J. (1990) 271, pp167-169.

The inventor has discovered that certain glutathione derivatives are effective inhibitors of the growth of a range of cancer cell types, and certain micro-organisms.

According to one aspect of the present invention there is provided a glutathione having the following structure:-

Where R1-R4 may be configured as shown in the following:

LABEL	R ₂	R_1	R ₄	R ₃	MW
CD4		B r	OMe	ОМе	638. 5
CD6	Сн₃со	в г	ОН	ОН	545
CD7	0 7 0	B r	ОМе	ОН	624

CD8		CH ₂ -COOEt	ОН	ОН	527
CD10			Оме	OMe	692
LABEL	R ₂	R ₁	R ₄	R ₃	MW
CD13	Н	NO 2	ОН	ОН	-

CD16	СНО	B r	ОН	ОН	504
CD17	Q =0	B r	NH ₂	ОН	576
CD19	0 0	NO 2	ОМе	ОН	621
CD20	0~0	NO2	ОМе	ОМе	635

CD42	N O 2	Оме	ОН	708
CD43	N O 2	ОМе	ОМе	722
CD46	-CH ₂ -CO ₂ Et	ОМе	ОМе	555

XCID: -WO__ 9809986A1 I_>

CD48 CH ₃ CO	ر ا ا ا	OMe	ОН	556	
-------------------------	------------------	-----	----	-----	--

CD7 has been found to be an effective inhibitor of Trypanosomiasis and in particular T. Brucei S247. This compound is also active against Malaria. One particular advantage of this compound is that it is effective at inhibiting the cell growth of the organism without being toxic to red blood cells.

10 CD13 is effective at the inhibition of growth of cancer cells, and in particular leukaemia, breast cancer or tumour cells.

According to another aspect of the invention there is

provided a glutathione having the following general formula:-

 $CH_2CH=C$

10 CH_3) $CH_2CH_2CH=C(CH_3)CH_2CH=C(CH_3)_2$

TRANS TRANS TRANS

S-(farnesyl)glutathione CD37

The aforementioned glutathione derivatives may each be provided in pharmaceutically acceptable compositions for delivery to the human or animal body.

In another aspect of the invention there is provided the use of any one of the foregoing compounds in the treatment of cancer, or parasitic cellular

20 infestation.

15

25

In particular, according to one embodiment of the present invention there is provided the use of CD13 and derivatives thereof in the treatment of cancer.

According to another embodiment there is provided the

use of CD19 and/or CD20 in the treatment of a infection by a parasitic micro-organism.

In particular Cd19 is highly effective at low concentrations against T. brucei (African sleeping sickness), while CD 20 has no toxicity to red blood cells. CD20 also has good activity against other parasites such as L.donovani (oriental sore, Kala-azar).

10

20

5

According to another aspect of the invention there is provided the use of CD7 against malaria.

According to yet another aspect of the invention there
is provided the use of any one of compounds CD26-CD37
against cancer.

According to another aspect of the invention there is provided the use of CD13, CD4, CD6 and CD 37 against cancer, and in particular breast cancer, and more particularly MCF7 cells.

Following is a description by way of example only of methods of putting the present invention into effect

25 and examples demonstrating the activity of compounds according to the present invnetion. The drawing is a

11

graphic representation of the results of the in vivo test described in example 1.

General method of production.

5

GSH + RX → GSR

Reduced glutathione (1g, 3.26mM) is dissolved in H₂0 (5ml) and 2M NaOH (3.3ml, 6.6mM) with stirring at room temperature and under a nitrogen atmosphere. Ethanol (5-15ml) is then added to the cloud point whereafter RX (for example, aryl halide, 3.5mM dissolved in ethanol) is added portion-wise over about 30 minutes. The reaction is left to stir for 20 hours under nitrogen.

If precipitation occurs during addition either more ethanol or more water is added to dissolve the material. At the end the reaction the acidity of the mixture is adjusted to p 3.5 with 2M HCl and the mixture chilled to effect precipitation. The precipitate is then filtered, washed with water, dried and recrystallized from Ethanol/H,0.

25 <u>Pharmaceutical</u> <u>activity:</u>

Example 1

Glutathione CD13 according to the present invention was tested in order to ascertain its inhibitory characteristics with respect to various cancer cell lines.

5

10

The compound was introduced to cell cultures and the concentration of cancer cells formed over a period of time was measured using standard techniques. The following table indicates the results for CD13 and another glutathione derivative "control" by way of comparison:-

	Leukaemia	lines	Tumour	Breast
			cells	Cancer
				cells
	WEH1 3B	K562	MAC 15A	MCF7
Compound	conc. μg/ml	conc. μg/ml	conc. μg/ml	conc. µg/ml
Control	36	54	>100	>100
CD13	3.4	2.5	0.43	0.48

15

20

CD13 was tested in vivo on rodents with MAC 15A S/C Tumours by giving them a 20 mg/Kg daily dose for five days. The treatment resulted in a reduction in tumour size and a 45% reduction in tumour volume after 4 days. The effect of CD 13 in the above test is shown

in the graph of the drawings.

Example 2

- The glutathione derivatives CD7 and CD 10 according to the present invention were tested for their activity against the parasites T.Brucei S247, L.donovani and T. cruzi.
- The following table shows the mean estimation of growth of T.Brucei S247 in a 72 hour incubation in the presence of a control glutathione derivative and CD10 in various concentrations:-

15	Compound	MIC @ concentration (µM)					
	Ť	30	10	3	1		
	control	++++	++++	++++	++++		
	CD7	0	++++	++++	++++		
	CD10	0	++++	++++	++++		

20

The forgoing table shows that Compounds CD7 and CD10 exhibit complete inhibition of the growth of T.Brucei S247 over the specified period at concentrations of 30 μM_{\odot}

The following table shows the results of the inhibition of the growth of L. donovani and T. Cruzi by compounds CD7, CD10 according to the present invention and "control" by way of comparison, at various concentrations.

Compound	% Inhi	bition	of	% Inhibition of			
	L.dono	ovani 0	conc.	T. Cruzi @ conc.			
	90 30 10			90	30	10	
	(µM)	(µM)	(Mu)	(µM)	(µM)	(µM)	
control	0	0	0	0	0	0	
CD7	0	0	0	0	0	0	
CD10	T	T	0	T	T	T/0	

10

5

The foregoing shows that compound CD10 shows good

15 activity in the inhibition of growth of both L.

donvani and T. cruzi.

Example 3:

20 Activity of compounds CD16-CD20 in Vitro.

Compound	% inhibition @ concn. (µ M)

٠. .

15

	30	10	3	1
CD16				
T.cruzi	11.0	0	0	0
CD17				
T.cruzi	7.0	4.0	0	0
CD19				
T.brucei	100	100	33.2	0
CD20				
L.donovani	13.5	1.9	0	0
T.brucei	100	64.5	0	0

10

5

Example 4:

Compound CD48 was tested for its activity against cancer, with the following results for a range of cancer types:

Cancer	Human Ovarian carcinoma	Human lung carcinoma	Human colon carcinoma	Human myclogenious leukaemia	Mouse lymphoid neoplasm
Designa tion	A2780	H-460	BE	K562	P388

	μМ	μМ	μМ	μМ	μМ
CD48	>50	>50	27.1	16.7	31.7

Example 5:

5

Compounds CD4, CD6, and CD 37 have also been found to be effective against various cancers, including breast cancer.

10 Example 6:

Compounds CD42 to 48 were tested for their activity against various parasitic infection agents and the results are shown in the following table:

15

Comp	% INHIBITION OF			%INHI	%INHIBITION OF L.			
ound	T.BRUCEI S247			DONOV	DONOVANI			
	30µM	10μΜ	ЗμМ	1μM	30µM	10μΜ	ЗμМ	lμM
CD42	100	100	0	0	1.2	0	0	0
CD43	100	100	0	0	T /O	0	0	0
CD44	100	100	100	63.2	T/	0	0	0
		 -		·	100			

WO 98/09986 PCT/GB97/02358

17

CD46	0	0	0	0	66.2	0	0	0
					3			

Key: T/0 = toxic to macrophages / parasites present.
T/100 = toxic to macrophages / no parasites
present.

100+ = 100% inhibition when sampling for
haemocytometer count but parasites visible in
wells of 96-well plate under inverted mic.

CLAIMS

1. A compound having the following general structure:-

5

and wherein R_1 to R_4 are according to any one of the following rows in the table:

LABEL	R ₂	R ₁	R ₄	R ₃
CD4		B 1	OMe	ОМе

CD6	сн _з со	81	ОН	ОН
CD7		B 1	OMe	ОН
CD8		СН ₂ -СООЕt	ОН	ОН

		-		
CD10			OMe	OMe
CD13	Н -	N C	ОН	ОН
CD16	СНО		ОН	ОН
CD17			NH ₂	ОН

100 mg/mg/

CD19	·/\"	H II 2	OMe	ОН
CD20		N 0 3	OMe	OMe
CD37	Ĥ	CH ₂ CH=C(CH ₃) CH ₂ CH ₂ CH=C(CH ₃)CH ₂ CH ₂ CH =C(CH ₃) ₂	ОН	ОН

SUBSTITUTE SHEET (RULE 26)

CD42	NO 2	OMe	ОĤ
CD43	N to Z	OMe	OMe
CD46	-CH ₂ -CO ₂ Et	ОМе	OMe

CD48 CH ₃ CO	OMe	ОН
-------------------------	-----	----

- A pharmaceutically acceptable composition for
 delivery to the human or animal body comprising a compound according to claim 1.
 - 3. A composition for the treatment of parasitic infection, which composition comprises one or more of CD7, CD19, CD20, CD42, CD43 or CD44 according to claim 1.
- A composition for the treatment of trypanosomiasis, and in particular infection by
 T. Brucei, which composition comprises one or more of CD7, CD19, CD20, CD42, CD43 and CD44 according to claim 1.
- 5. A composition as for the treatment of

 Leishmaniasis, and in particular infection by L.

 donovani, which composition comprises one or

 more of CD20, CD42, CD43, CD44 and CD46 according

10


15

24

to claim 1.

- 6. A composition for the treatment of cancer, which composition comprises one or more of CD4, CD6, CD13 and CD37 according to claim 1.
- 7. A method of treating a diseased human or animal comprising administering a pharmologically effective amount of a composition as claimed in claim 2.
- 8. A method of treating parasitic infection of a human or animal comprising administering a pharmologically effective amount of a composition as claimed in any of claims 3,4 and 5.
- A method of treating cancer in a human or animal comprising administering a pharmologically effective amount of a composition as claimed in claim 6.

MAC15A S/C TUMOURS TREATED WITH CD13 DAILY



INTERNATIONAL SEARCH REPORT

Intern. .nai Application No PCT/GB 97/02358

		Į PL	CT/GB 97/02358
IPC 6	FICATION OF SUBJECT MATTER C07K5/02 A61K38/05		
According to	o International Patent Classification (IPC) or to both national clas	sification and IPC	
3. FIELDS	SEARCHED		
Minimum do IPC 6	ocumentation searched (classification system followed by classi CO7 K	ication symbols)	
Documentat	tion searched other than minimum documentation to the extent t	nat such documents are included in	n the fields searched
Electronic d	late base consulted during the international search (name of dal	a base and, where practical, searc	oh terms used)
C. DOCUM	ENTS CONSIDERED TO BE RELEVANT		
Category *	Ctation of document, with indication, where appropriate, of the	e relevant passages	Relevant to claim No.
X	R.VINC ET AL.: "Studies on the of Glyoxalase I by S-Substitut Gluthationes"		1,6-9
	J.MED.CHEM., vol. 14, no. 5, 1971,		
,	page 402-4 XP002044716 see paragraph 1; example 39; t		
X	lyase) by N-acylated S-blocked derivatives as a probe for rol	lian glyoxalase I (lactoylglutathione) by N-acylated S-blocked glutathione atives as a probe for role of the e of gluthathione in glyoxalase I	
	vol. 86, no. 1, 1986, pages 160-8, XP002044717 see tables 12,5		
		,	
	<u> </u>	-/	
X Furti	her documents are listed in the continuation of box C,	X Patent family memb	ers are listed in annex.
° Special ca	stegories of cited documents :	'T' later document nublisher	after the international filing date
consid	ent defining the general state of the art which is not dered to be of particular relevance document but published on or after the international	or priority date and not in orted to understand the invention	in conflict with the application but principle or theory underlying the
filing d "L" docume which		cannot be considered n involve an inventive ste "Y" document of particular re	levance; the claimed invention ovel or cannot be considered to p when the document is taken alone devance; the claimed invention
	ent referring to an oral disclosure, use, exhibition or	document is combined a menta, such combination	o involve an inventive latep when the with one or more other such docu- in being obvious to a person skilled
P° docume	ent published prior to the international filing date but han the priority date claimed	in the art. "&" document member of the	same patent family
	actual completion of the international search 7 October 1997	Date of mailing of the int	emational search report 1 3. 11, 97
	mailing address of the ISA	Authorized officer	
	European Patent Office, P.B. 5818 Patentiaan 2 NL - 2280 HV Rijewijk Tel: (+31-70) 340-2040;Tx: 31 651 epo nt; Fax: (+31-70) 340-3016	Deffner, (C-A

Form PCT/ISA/210 (second sheet) (July 1992)

INTERNATIONAL SEARCH REPORT

Intern. _nal Application No PCT/GB 97/02358

10000	DOCUMENTS CONCINEDED TO BE DELEVANT	PCT/GB 97/02358	
(Continua tegory °	ation) DOCUMENTS CONSIDERED TO BE RELEVANT Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.	
wyury *	онавон он посиняли, эти иносания, этим африсунали, он не темуал раззадех	neevant to caum No.	
	A.AL-TIMARI ET AL.: "Inhibition of glutathione derivatives of bovine liver glyoxalase II (Hydroxyacylgluthatione hydrolase) as a probe of the N- and S-sites for substrate binding" BIOCHIM.BIOPHYS.ACTA, vol. 870, no. 2, 1986, page 219-25 XP002044718 see example 2; table I	1,6-9	
A	S.J.NORTON ET AL.: "Inhibitors and inhibition of mammmalian glyoxalase II activity" BIOCHEM.SOC.TRNS., vol. 21, no. 2, 1993, pages 545-9, XP002044719		
A	NORTON S.J. ET AL.: "Glyoxalase and Glyoxalase II from Aloe Vera: Purification, Characterization and Comparision with Animal Glyoxalases" BIOCHEM.INT, vol. 22, no. 3, 1990, pages 411-18, XP002044720		
A	AC. ELIA ET AL.: "N,S-Bis-Fluorenylmethoxycarconylglutathio ne: A New, Very Potent Inhibitor of Mammalian Glyoxalase II" BIOCHIM.MOL.BIOL.INT., vol. 35, no. 4, 1995, pages 763-71, XP002044721		
Α	WO 95 08563 A (TERRAPIN TECH INC) 30 March 1995		
	·		
**	.	-	

INTERNATIONAL SEARCH REPORT

Information on patent family members

PCT/GB 97/02358

Patent document	Publication	Patent family member(s)	Publication
cited in search report	date		date
WO 9508563 A	30-03-95	US 5599903 A AU 7842194 A CA 2171453 A EP 0720620 A JP 9506336 T US 5556942 A	04-02-97 10-04-95 30-03-95 10-07-96 24-06-97 17-09-96

Form PCT/ISA/210 (patent lamsly annex) (July 1992)